

RIB1213

A.oryzae var. *brunneus* Murakami



Colony on CD plate



Colony on koji extract plate

Mycological characters

| Color of old slant culture | Diameter (mm) of colony at 34 C | Reverse side of Colony | | Sclerotia |
|-----------------------------------|--|--------------------------|--------------------------------------|-------------------------|
| | | Wrinkling | Color | |
| 1 | 4 | 4 | 2 | 1 |
| Roughness of stalk | Length of stalk (mm) | Diameter of vesicles (μ) | Roughness of conidia | Diameter of conidia (μ) |
| 6 | 6 | 2 | 3 | 5 |
| Ratio of biseriata sterigmata (%) | Production | | | |
| | Kojic acid | Pigments | Aflatoxins | Amylase |
| 1 | 3 | 3 | 1 | 4 |
| Coloration of hydroquinone | Reduction of methylene blue | Browning of rice koji | Pink color of conidia in meidum with | |
| | | | Anisic acid | Anisaldehyde |
| 5 | 1 | 5 | 2 | 4 |
| Note | IAM 2745 <i>A. oryzae</i> S. et Y, (1969)-ACTU. sakaguchi A-4-1 ITS type A1a, AFHC type <i>A. oryzae</i> group2 (Tominaga et. al, AEM, 72, 484 (2006), Kiyota et. al, JBB, 111, 512 (2011)) | | | |